

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Neurotoxicity of the Parkinson's Disease-Associated Pesticide Ziram Is Synuclein-Dependent in Zebrafish Embryos

Aaron Lulla, Lisa Barnhill, Gal Bitan, Magdalena I. Ivanova, Binh Nguyen, Kelley O'Donnell, Mark C. Stahl, Chase Yamashiro, Frank-Gerrit Klärner, Thomas Schrader, Alvaro Sagasti, and Jeff M. Bronstein

Table of Contents

Figure S1: ZF $\gamma 1$ antibody is specific for ZF. Using denaturing conditions/SDS PAGE, a band for $\gamma 1$ was detected at 17 kDa for ZF adult brain (ZF) and purified ZF $\gamma 1$ (a). No ZF $\gamma 1$ signal was detected for mouse brain (MB). Using non-denaturing conditions/Native PAGE, a major band for ZF $\gamma 1$ was detected at 480 kDa and a minor band at 242 kDa (b). Peptide preincubated with the ZF $\gamma 1$ antibody is shown for SDS PAGE. No band for ZF $\gamma 1$ was detected after preincubation of $\gamma 1$ antibody with $\gamma 1$ peptide (c). ZF $\gamma 1$ MO reduced protein levels of ZF $\gamma 1$ as compared to scramble MO as determined by SDS PAGE (d).

Figure S2: Ziram is toxic in a concentration-dependent manner and causes notochord distortion. A significant change in toxicity was observed for embryos treated (24hpf) with 100nM and 1 μ M ziram by day 7 ($n = 50$, $p < .0001$) (a). ZF embryos treated with 50 nM ziram at 5 hpf, were found to have a shorter body axis, pericardial edema, and notochord distortion (b) as compared to vehicle treated ZF (c). $p < .0001$ Log-Rank test.

Figure S3: Ziram treatment results in reduction of TH-1 levels. Using Western blot analysis, the effect of ziram (50nM) on TH-1 levels was investigated. A 63% reduction in TH-1 was observed for embryos treated with 50nM ziram vs. controls (n=4; *p-value = .03 using two-tailed T test). Bars represent standard error of the mean.




Figure S1: ZF $\gamma 1$ antibody is specific for ZF. Using denaturing conditions/SDS PAGE, a band for $\gamma 1$ was detected at 17 kDa for ZF adult brain (ZF) and purified ZF $\gamma 1$ (a). No ZF $\gamma 1$ signal was detected for mouse brain (MB). Using non-denaturing conditions/Native PAGE, a major band for ZF $\gamma 1$ was detected at 480 kDa and a minor band at 242 kDa (b). Peptide preincubated with the ZF $\gamma 1$ antibody is shown for SDS PAGE. No band for ZF $\gamma 1$ was detected after preincubation of $\gamma 1$ antibody with $\gamma 1$ peptide (c). ZF $\gamma 1$ MO reduced protein levels of ZF $\gamma 1$ as compared to scramble MO as determined by SDS PAGE (d).





Figure S2: Ziram is toxic in a concentration-dependent manner and causes notochord distortion. A significant change in toxicity was observed for embryos treated (24hpf) with 100nM and 1µM ziram by day 7 ($n = 50$, $p < .0001$) (a). ZF embryos treated with 50 nM ziram at 5 hpf, were found to have a shorter body axis, pericardial edema, and notochord distortion (b) as compared to vehicle treated ZF (c). $p < .0001$ Log-Rank test.




Figure S3. Ziram treatment results in reduction of TH-1 levels. Using Western blot analysis, the effect of ziram (50nM) on TH-1 levels was investigated. A 63% reduction in TH-1 was observed for embryos treated with 50nM ziram vs. controls (n=4; *p-value = .03 using two-tailed T test). Bars represent standard error of the mean.